

Coastwide Nutria Control Program 2009-2010

**Nutria Harvest and Distribution 2009-2010
and
A Survey of Nutria Herbivory Damage in Coastal
Louisiana in 2010**



**Conducted by:
Coastal and Nongame Resources
Louisiana Department of Wildlife and Fisheries**

**As part of the Coastwide Nutria Control Program*
CWPPRA Project (LA-03b)**

Submitted by: Jillian Jordan and Edmond Mouton June 30, 2010

*Funded by Coastal Wetlands, Planning, Protection, and Restoration Act through the Natural Resources Conservation Service and the Office of Coastal Protection and Restoration.

TABLE OF CONTENTS

	Page
Section 1	Nutria Harvest Distribution 2009-2010----- 4-12
Section 2	A Survey of Nutria Herbivory Damage in Coastal Louisiana in 2010----- 13-20
Section 3	Summary----- 20-22
 Figures	
	Figure 1 (Annual Harvest and Avg. Price of Nutria) ----- 5
	Figure 2 (Range of Nutria Harvest) ----- 9
	Figure 3 (Nutria Harvest per Month) ----- 9
	Figure 4 (Harvest by Marsh Type) ----- 10
	Figure 5 (Method of Take) ----- 10
	Figure 6 (Method of Take by Marsh Type) ----- 11
	Figure 7 (Harvest by Parish) ----- 12
	Figure 8 (Damage Type) ----- 16
	Figure 9 (Percentage of Damaged Acres per Parish) ----- 17
	Figure 10 (Damaged Acres by Marsh Type) ----- 17
	Figure 11 (Nutria Relative Abundance Rating) ----- 18
	Figure 12 (Vegetative Damage Rating) ----- 18
	Figure 13 (Age of Damage and Condition Rating) ----- 19
	Figure 14 (Prediction of Recovery) ----- 19
	Figure 15 (CNCP Progress) ----- 21
 Tables	
	Table 1 (Carcass Use) -----11
	Table 2 (Muskrat Damage) -----20
	Table 3 (Three Years Prior to CNCP) ----- 21
	Table 4 (First Eight Years of the CNCP) ----- 21
	Table 5 (Nutria Harvest by Parish Seasons 1-8) ----- 24-25
	Table 6 (Method of Take by Parish Seasons 1-8) ----- 26-27
	Table 7 (Status and Number of Nutria Herbivory Sites) ----- 28
	Table 8 (Number of Nutria Damage Sites and Acres) ----- 29-30
	Table 9 (Number of Damage Sites and Acres by Marsh Type) ----- 31
	Table 10 (Nutria Relative Abundance Rating) -----32-33
	Table 11 (Vegetative Damage Rating) ----- 34-35
	Table 12 (Age of Damage and Condition Rating) ----- 36-37
	Table 13 (Prediction of Recovery) ----- 38-39
	Table 14 (2009 Nutria Veg. Damage Survey Results) ----- 41
	Table 15 (2010 Nutria Veg. Damage Survey Results) ----- 44

Appendices

Appendix A -----	23-39
- Season Comparison Data	
Appendix B -----	40-42
- 2009 Nutria Veg. Damage Sites	
- 2009-2010 Harvest and 2010 Damage Sites Map	
Appendix C -----	43-46
- 2010 Veg. Damage Survey Results	

Section 1

NUTRIA HARVEST DISTRIBUTION 2009-2010

Introduction

Since 2001, annual coast wide aerial surveys assessing herbivory in Louisiana have documented approximately 26,273 acres of marsh converted to open water due to nutria vegetative damage. (This acreage is actual observed acreage multiplied by a constant to account for land not seen from the transects.) This loss of marsh in Louisiana is devastating to the people that depend on the marsh for their livelihood as well as people that use it for recreation. It is vital to the people of Louisiana to protect the wetlands from destruction whenever possible. In order to remove the threat of land loss due to nutria, the Coastwide Nutria Control Program was developed.

The nutria (*Myocastor coypus*) is a large semi-aquatic rodent indigenous to South America. The first introduction of nutria to North America occurred in California in 1899; however it was not until the 1930's that additional animals were introduced in seven other states. These importations, primarily for fur farming, failed during the Second World War as a result of poor pelt prices and poor reproductive success. After the failures of these fur farms, nutria were released into the wild. Sixteen states now have feral populations of nutria.

The Gulf Coast nutria population originated in Louisiana in the 1930's from escapes and possible releases from nutria farms. Populations first became established in the western coastal portion of the state and then later spread to the east through natural expansion coupled with stocking. During the mid-1950s muskrat populations were declining, nutria had little fur value, and serious damage was occurring in rice fields in southwestern Louisiana and sugarcane fields in southeastern Louisiana; farmers complained about damage to crops and levee systems, while muskrat trappers blamed the nutria for declining numbers of muskrats. In 1958, the Louisiana Legislature placed the nutria on the list of unprotected wildlife and created a \$0.25 bounty on every nutria killed in 16 south Louisiana parishes, but funds were never appropriated.

Research efforts were initiated by the federal government in the southeastern sugarcane region of the state to determine what control techniques might be successful. This research conducted by the U.S. Fish and Wildlife Service during the 1960's examined movements in relation to sugarcane damage and recommended shooting, trapping, and poisoning in agricultural areas. Ted O'Neil, Chief of the Fur and Refuge Division, Louisiana Department of Wildlife and Fisheries (LDWF), believed that the problem could only be solved through the development of a market for nutria pelts. A market for nutria developed slowly during the early 1960's and by 1962 over 1 million pelts were being utilized annually in the German fur trade. The nutria became the backbone of the Louisiana fur industry for the next 20 years, surpassing the muskrat in 1962 in total numbers harvested. In 1965, the state legislature returned the nutria to the protected list. As fur prices showed a slow rise during most of the 1970's and early 1980's, the harvest averaged 1.5 million pelts and complaints from agricultural interest became uncommon. From 1971 through 1981 the average annual value of the nutria harvest to the coastal trappers was \$8.1 million. The nutria harvest in Louisiana from 1962 until 1982 remained over 1 million annually. The harvest peaked in 1976 at 1.8 million pelts worth \$15.7 million to coastal trappers (Figure 1).

The nutria market began to change during the early 1980's. In 1981-1982, the nutria harvest dropped slightly below 1 million.

This declining harvest continued for two more seasons; then in the 1984-1985 season, the harvest jumped back up to 1.2 million. During the 1980-1981 season, the average price paid for nutria was \$8.19. During the 1981-1982 season, the price dropped to \$4.36 and then in 1982-1983, the price dropped to \$2.64. Between the 1983-1984 season and the 1986-1987 season, prices fluctuated between \$3.00 and \$4.00. Then in 1987-1988 and again in 1988-1989 prices continued to fall (Figure 1). From 1982 through 1992 the average annual value of the nutria harvest was only \$2.2 million. Between 1988-1989 and 1995-1996 the number of nutria harvested annually remained below 300,000 and prices remained at or below a \$3.00 average.

Due to a strong demand for nutria pelts in Russia in both 1996-1997 and in 1997-1998, 327,286 nutria were harvested at an average price of \$4.13 and 359,232 nutria were harvested at an average price of \$5.17 during those seasons respectively. In September 1998, the collapse of the Russian economy and general instability in the Far East economies weakened the demand for most wild furs including nutria. The demand for nutria pelts in Russia declined quickly due to the devaluation of the Russian ruble. During the 1998-1999 trapping season, pelt values fell to \$2.69 and harvest decreased to only 114,646, less than one-third of the previous year. During the 1999-2000 trapping season there was virtually no demand for nutria pelts. The harvest decreased to 20,110 nutria. This was, by far, the lowest nutria harvest on record since the mid-1950s. The number of nutria harvested in 2000-2001 trapping season increased to 29,544. The value of nutria pelts decreased to \$1.75 during the 2001-2002 season, prompting another decrease in harvest to 24,683 nutria.





Figure 1



During the strong market period for nutria pelts, there were no reports of wetland damage caused by nutria. However, before the market developed and after the market declined, reports of marsh vegetation damage from land managers became common. Such complaints began in 1987 and became more frequent during the early 1990's. In response, the Fur and Refuge Division of the Louisiana Department of Wildlife and Fisheries (LDWF) initiated limited aerial survey flights, particularly in southeastern Louisiana. Survey flights of Barataria and Terrebonne basins were conducted during the 1990's, with initial support from Barataria-Terrebonne National Estuary Program (BTNEP) and later support from Coastal Wetlands Planning, Protection and Restoration Act (CWPPRA). From 1993 to 1996 these flights showed acres of damage increasing from approximately 45,000 to 80,000 acres within the basins. The first CWPPRA funded coast wide survey, conducted in 1998, showed herbivory damage areas totaling approximately 90,000 acres. By 1999 this coast wide damage had increased to nearly 105,000 acres.

This rapid and dramatic increase in damaged acres prompted LDWF to pursue funding for the Coastwide Nutria Control Program (CNCP) in January 2002.

The project is funded by the CWPPRA through the Natural Resources Conservation Service (NRCS) and the Office of Coastal Protection and Restoration (OCPR) with the LDWF as the lead implementing agency. Task one requires LDWF to conduct an annual aerial survey to evaluate the herbivory damage caused by nutria. Task two of the OCPR and LDWF Interagency Agreement No. 2511-02-29 for the CNCP requires LDWF to conduct general project operation and administration. LDWF is required to 1) conduct and review the registration of participants in the CNCP; 2) establish collection stations across coastal Louisiana; 3) count valid nutria tails and present participants with a receipt/voucher; 4) deliver tails to an approved disposal facility and receive documentation that ensures the nutria will be properly disposed of and shall not leave the facility; and 5) process and maintain records regarding participants, number and location where tails were collected. Task 3 requires LDWF to provide incentive payments to program participants and task 4 requires LDWF to provide a report regarding the distribution of the harvest by township.

The program area is coastal Louisiana bounded to the north by Interstate-10 from the Texas state line to Baton Rouge, Interstate-12 from Baton Rouge to Slidell, and Interstate-10 from Slidell to the Mississippi state line. The project goal is to significantly reduce damage to coastal wetlands attributable to nutria herbivory by removing 400,000 nutria annually. This project goal is consistent with the Coast 2050 common strategy of controlling herbivory damage to wetlands. The method chosen for the program is an incentive payment to registered trappers/hunters for each nutria tail delivered to established collection centers. Initially, registered participants were given \$4.00 per nutria tail. To encourage participation, the payment was increased to \$5.00 per tail in the 2006-2007 season.

This section reports on the Nutria Harvest Distribution for 2009-2010.

Methods

The application for participation in the Coastwide Nutria Control Program (CNCP) was developed in July 2002 but was modified in June 2003 to obtain better information about the location of nutria harvest. It was made available through the LDWF offices and website, as well as LSU Cooperative Extension offices. In order for a participant to be qualified, the individual must complete the application, obtain written permission from a landowner or land manager with property in the program area, complete a W-9 tax form and provide LDWF with a complete legal description of the property to be hunted or trapped. A map outlining the property boundaries was an added requirement of participants beginning with the 2003-2004 season. Once an applicant was accepted, the participant was mailed information on the program's regulations, collection sites for nutria tails, contact information and a CNCP registration card.

Coastal Environments Inc. (CEI) was selected as the contractor to develop and maintain the program database, collect nutria tails, and distribute incentive payment checks to participants for tail harvests. The contract with CEI, which began with the 2002-2003 season, was extended to include the 2003-2004 through 2006-2007, with the option to renew for 3 years there after. CEI's first renewal season was (2007-2008), the second renewal season was (2008-2009), and they just completed their third renewal season (2009-2010). Tail collection sites were originally established at Rockefeller Refuge, Abbeville, Berwick (Morgan City), Houma, Luling and Slidell.

Collections were made once a week at each site, except for Rockefeller Refuge, Abbeville and Slidell, where collections were made by appointment only, due to low numbers of participants in those areas.

Louisiana's open trapping season began on November 20, 2009, and nutria tail collections began a week later. Collections were made utilizing a 16 foot by 8 foot trailer containing a freezer, sorting table and desk. A participant reported to a collection site, presented his nutria control program registration card and presented his tails to a CEI representative.



One CEI representative conducted an exact count of the nutria tails, which was then verified with the participant to ensure they were in agreement. At that time, the counted tails were placed into a plastic garbage bag labeled with the participant's CNCP registration number and the number of tails contained in that bag. Another CEI representative filled out a voucher for the number of tails delivered, checking to make sure the mailing address of the participant was correct. The participant was asked to provide the following information: 1) the method of taking the nutria, 2) the method in which the nutria carcass was used or abandoned, and 3) the month or months in which the nutria were harvested. When complete, the voucher was signed by the participant who would also indicate on a detailed map of their lease the location or locations where the nutria were harvested. The CEI representative recorded township and range of harvest, number of nutria harvested, and the transaction number on the map. One copy of the voucher was given to the participant, while one copy was retained by the CEI representative.



The information on the voucher was entered into a laptop computer and transferred electronically to the CEI main offices via an FTP site for analysis and quality control.

The data transfer occurred at the end of each collection day. Collected tails were transported to the BFI waste storage facility in Sorrento, Louisiana, at the end of each collection day or multiple times a day if necessary. The CEI representative checked in at a guard station where the vehicle containing the tails was weighed. The vehicle was also weighed when exiting the disposal site in order to calculate the exact amount of waste deposited at the facility. The tails were deposited into a biohazard waste pit under supervision of a BFI employee. The number of bags disposed, as well as weight deposited, was recorded on a receipt given to the CEI representative. Copies of the receipts for all disposals made were supplied to LDWF.

At the end of the collection week, the maps were transported to CEI's office in Baton Rouge. At this time QA/QC of the data transferred for the entire week took place. The trapped/hunted areas that were outlined on the lease maps were digitized into Arc Map GIS 9.2. CEI sent a weekly report to LDWF detailing each transaction, including a digitized map of that week's trapped/hunted areas. Each Monday morning, after receiving a weekly report and bill, LDWF sent a payment to CEI for the amount of tails collected and services rendered. CEI in turn sent participants checks through the mail for the amount of tails turned in. Louisiana's open trapping season ended on March 31, 2010, and nutria tail collections continued for one week into April. After the conclusion of the season, CEI provided LDWF with all the transaction information for the entire season from November to March. This final report contains information recorded on the vouchers, the digitized trapped/hunted area, the nutria control program database and an Arc Map 9.2 project map with related information.

Results and Discussion

Participant Totals

A total of 445,963 nutria tails, worth \$2,229,815 in incentive payments, were collected from 306 participants in the 2009-2010 season. Approximately 42% of these participants turned in 800 or more tails (Figure 2.)




Figure 2

Harvest by Month

The 2009-2010 trapping season began November 20th, 2009 and continued through March 31st, 2010. One hundred thirty thousand two hundred and six (130,206) tails were collected in the month of January making it the most active month of the season (Figure 3.)




Figure 3

Harvest by Marsh Type

Harvest data were classified by marsh type, which includes: fresh marsh, intermediate marsh, brackish marsh, salt marsh and other. The category “other” includes swamp, mixed forest, open water and agriculture land types.

In the 2009-2010 season, 46% of the nutria harvested fell into the “Fresh Marsh” category, followed by 34% being harvested from the “Other” (Figure 4.).




Figure 4

Method of Take

During collection transactions, program participants indicated their method of take: trapped, shot with rifle, or shot with shotgun.

The predominant method used in the 2009-2010 season was shooting with a rifle (Figure 5.)




Figure 5

While shooting with a rifle was the most popular method of taking nutria in fresh marsh and salt marsh, trapping was the most utilized method in brackish (Figure 6.)




Figure 6

Carcass Use

Use of nutria carcasses, was recorded for each participant transaction. For the purpose of this survey, use categories include: 1) harvested for meat and/or 2) harvested for fur (Table 1.)

MARSH TYPE	Fur	Meat	Abandon Buried	Abandon Vegetation	Abandon Water
Fresh	2,403	2,724	145,473	35,219	22,453
Intermediate	314	331	47,667	13,944	10,734
Brackish	86	123	9,171	1,821	609
Salt	36	50	3,913	0	177
Other	319	556	121,973	23,255	5,386
Total	3,158	3,784	328,197	74,239	39,359

Table 1

Overall, almost 2% of the nutria harvested was utilized for meat and/or fur. This is down 81% from the utilization last season. The remaining 98% were disposed of by approved methods, categories include: 1) buried carcasses, 2) placed in heavy overhead vegetation, or 3) placed in water (Table 1.)

All interested participants were supplied a fur buyer/fur dealer list to encourage the use of animals for the fur and meat, and interested fur buyers/dealers were supplied with a list of program participants.

Harvest by Parish

Nineteen parishes were represented in the 2009-2010 season of the Coastwide Nutria Control Program, with nutria harvests ranging from 267 to 106,226. Terrebonne Parish reported the highest number of tails with 106,226 followed by Plaquemines and St. Mary Parish with 69,294 and 67,631 respectively (Figure 7).




Figure 7

Section 2

A SURVEY OF NUTRIA HERBIVORY DAMAGE IN COASTAL LOUISIANA IN 2010

Introduction

Herbivory damage was noticed in the late 1980s by landowners and land managers when the price of fur dropped and the harvest of nutria all but ceased. The LDWF was contacted to investigate the problem. The first region wide aerial survey became possible because of the interest and concern of many state and federal agencies, coastal land companies and, in particular, funding provided by BTNEP. The objectives of the aerial survey were to: (1) determine the distribution of damage along the transect lines as an index of region wide damage, (2) determine the severity of damage as classified according to a vegetative damage rating, (3) determine the abundance of nutria by the nutria relative abundance rating (4) determine the species of vegetation being impacted and (5) determine the status of recovery of selected damaged areas (Linscombe and Kinler 1997).

Helicopter surveys were flown in May and December 1993 and again in March and April 1996 across the Barataria and Terrebonne Basins. During the December 1993 survey, 90 damaged sites were observed with more than 15,000 acres of marsh impacted along the transects and an estimated 60,000 acres across the study area. In 1996, a total of 157 sites were observed. The damage observed along the transect lines increased to 20,642 acres, and an extrapolated acreage of 77,408 acres across the study area. (The extrapolated coast wide estimate is derived by multiplying the observed acres by 3.75 to account for area not visible from the transect lines.) All of the 1993 sites were evaluated again in 1996, but only 9% showed any recovery. Clearly, the trend identified was a continued increase in both the number of sites and the extent of nutria damage in the Barataria and Terrebonne Basins.

In 1998, the first coast wide nutria herbivory survey was flown, as part of the Nutria Harvest and Wetland Demonstration Program (LA-03a). A total of 23,960 acres of damaged wetlands were located at 170 sites along the survey transects, with an extrapolated coast wide estimate of 89,850 acres. In 1999, the damage increased to 27,356 acres located at 150 sites, with an extrapolated coast wide estimate of 102,585 acres. In 2000, the damage slightly decreased to 25,939 acres located at 132 sites, with an extrapolated coast wide estimate of 97,271 acres. In 2001, the damage decreased to 22,139 acres located at 124 sites, with an extrapolated coast wide estimate of 83,021 acres. In the 2002 survey, the first survey funded as part of the CNCP and the survey which preceded implementation of the CNCP incentive payments, the damage decreased again, but only slightly to 21,185 acres located at 94 sites, with an extrapolated coast wide estimate of 79,444 acres. During the 2003 survey, a total of 84 sites had some level of vegetative damage and covered a total of 21,888 acres, with an extrapolated coast wide estimate of 82,080 acres. In summary, the coast wide estimates of nutria herbivory damage prior to implementation of the CNCP incentive payments (from 1998 to 2003) ranged from 79,444 to 102,585 acres.

Vegetative damage caused by nutria has been documented in at least 11 Coastal Wetlands Planning Protection and Restoration Act (CWPPRA) project sites in the Barataria and Terrebonne Basins. Nutria herbivory is only one of many factors causing wetlands loss, but the additional stress placed on the plants by nutria herbivory may be very significant in CWPPRA projects sites and throughout coastal Louisiana.

The previous extrapolated estimates of 79,444 to 102,585 acres of marsh damaged was conservative because only the worst sites (most obvious) can be detected from aerial surveys; the actual number of acres being impacted was certainly higher. When vegetation is removed from the surface of the marsh, as a result of over grazing by nutria, the very fragile organic soils are exposed to erosion through tidal action and/or storms. If damaged areas do not revegetate quickly, they may become open water as tidal scour removes soil and thus lowers elevation. This is evident as the damaged sites that converted to open water over the last five years have been in the intermediate and brackish marsh types. Frequently the plant's root systems are also damaged, making recovery through vegetative regeneration very slow.

In an effort to create an incentive for trappers and hunters, the CNCP was implemented. Task number 1 of the LDNR and LDWF Interagency Agreement No. 2511-02-29 for the CNCP requires LDWF to conduct annual coast wide aerial surveys during spring/summer to document the current year impact of nutria herbivory. Survey techniques followed Linscombe and Kinler (1997), and CNCP funded surveys, have been conducted each spring from 2003 to 2010. Results were analyzed and the numbers of acres impacted or recovered were determined.


This section reports on the 2010 Coastwide Nutria Herbivory Survey.

Methods



The 2010 coast wide nutria herbivory survey was conducted May 5th- 8th and May 10th-14th. North-South transects were flown throughout the fresh, intermediate and brackish marshes of coastal Louisiana. Under normal circumstances, a total of 155 transects (covering 2,354.7 miles) are surveyed for damage. The transects were spaced approximately 1.8 miles apart, starting at the swamp-marsh interface and continuing south to the beginning of the salt marsh. Due to low nutria population density, salt marsh habitat was not included in the survey. Depending upon visibility and vegetative conditions, an altitude of 300-400 feet was considered optimum. At this altitude, vegetative damage was identifiable and allowed for a survey transect width of about 1/4 mile on each side of the helicopter. Flight speed was approximately 60 mph. Two observers were used to conduct the survey, each positioned on opposite sides of the helicopter. In addition to locating vegetative damage, one observer navigated along the transect line and the other observer recorded all pertinent data.

However, due to the BP oil spill in the Gulf of Mexico a Temporary Flight Restriction (TFR) was implemented at the mouth of the Mississippi River. With high air traffic and safety concerns, a small area was excluded from the survey (See Map Below). Even though this area was not flown, it is not thought to interfere with the survey data. Since the aerial vegetative damage survey started in 1997, there has been no damage recorded along the excluded transects.



When vegetative damage was identified, the following information was recorded.

1) Location of each site was determined by recording latitude and longitude utilizing GPS equipment. A real time differential corrected (WAAS Enabled) GPS (Garmin GPSmap 296) was utilized to allow for accurate location of damaged sites. The software used was DNRGarmin (written by Minnesota DNR) operating in ArcView 9.2. The size of each damage site was recorded by logging polygons using stream digitizing with the GPS equipment.

2) The abundance of nutria sign was placed in one of the following nutria relative abundance rating (NRAR) categories: **no nutria sign visible (0)**, **nutria sign visible (1)**, **abundant feeding (2)**, **heavy feeding (3)**.

3) The extent of damage to the vegetation was placed in one of the following vegetative damage rating categories: **no vegetative damage (0)**; **minor vegetative damage (1)** which is defined as a site containing feeding holes, thinning vegetation and some visible soil; **moderate vegetative damage (2)** which is defined as a site that has large areas of exposed soil and covers less than 50% of the site; **severe vegetative damage (3)** which is defined as a site that has more than 50% of the soil exposed; or **converted to open water (4)**.

4) The dominant plant species were identified and recorded for damaged areas, recovering areas and in the adjacent areas.

5) The age of damage and condition is determined by considering feeding activity and vegetation condition. The age of damage and condition was placed in one of the following categories: **recovered (0), old recovering (1), old not recovering (2), recent recovering (3), recent not recovering (4) or current (occurring now)(5).**

6) The prediction of vegetative recovery is made considering feeding activity, age of damage and the extent of damage. The prediction of vegetative recovery by the end of 2010 was characterized by one of the following categories: **no recovery (0), full recovery (1), partial recovery (2) or increased damage (3).**

7) The number of nutria observed at each site was recorded.

In addition to searching for new damaged sites, all previously identified damaged sites were revisited to assess extent and duration of damage or to characterize recovery. All data were entered into a computer for compilation. Damaged site locations are provided on the attached herbivory map and a data summary in Appendix B.

Results and Discussion

There were 22 sites included in the 2010 vegetative damage survey. Of the 22 sites, 20 were previously classified as damaged sites in the 2009 survey, 1 converted from recovered in 2009 to damaged in 2010, and there was 1 new site located. Nine of the nutria damage sites and 1 of the muskrat/storm sites from 2009 have completely recovered. One site has converted to being all hog damage; whereas in 2009 it was ½ nutria and ½ hog damage. The remaining 11 sites are classified as nutria damage sites (Figure 8.).




Figure 8

Nutria Damage

The following discussion details the 11 sites that had nutria damage (Appendix A). A total of 2,260 acres along transects (extrapolated to be 8,475 acres coast wide) in 2010, were impacted by

nutria feeding activity. This represents approximately a 58% decrease in acres impacted by nutria in 2009 (5,422 acres, extrapolated 20,333 acres coast wide.)

Damage by Parish

Terrebonne parish experienced almost all of the damaged acres in 2010 (Figure 9).




Figure 9

Damage by Marsh Type

Marsh type was recorded for each damage site, as well as the type of vegetation based on the Linscombe and Chabreck 2001 survey (Figure 10.). All 11 nutria damage sites were within fresh marsh during the 2010 survey.




Figure 10

Fresh marsh continued to be the most affected by nutria herbivory (~100%). The typical vegetation impacted in fresh marsh was *Eleocharis* spp. and *Hydrocotyle* spp.

Nutria Relative Abundance Rating

A nutria relative abundance rating (NRAR) was used to quantify the abundance of nutria at each site. Categories include: (0) no nutria sign visible, (1) nutria sign visible, (2) abundant feeding sign, and (3) heavy feeding sign; sites converted to open water are not given a NRAR (Figure 11.)




Figure 11

Vegetative Damage Rating

Vegetative damage was also evaluated at each site. A rating system was developed in order to quantify nutria vegetative damage. The vegetative damage rating (VDR) has five categories: (0) no vegetative damage, (1) minor vegetative damage, (2) moderate vegetative damage, (3) severe vegetative damage, (4) converted to open water (Figure 12.)




Figure 12

Age of Damage Rating

Categories for the age of damage and condition rating include: (1) current damage, (2) recent damage-recovering, (3) recent damage not recovering, (4) old damage-recovering, (5) old damage-not recovering, and (0) recovered (Figure 13.)




Figure 13

Prediction of Recovery

For each site with current damage, the degree of recovery by the end of the 2010 growing season was predicted. These categories include: (1) full recovery, (2) partial recovery, (3) increased damage and (4) no recovery predicted (Figure 14.)




Figure 14

Muskrat Damage

During the 2010 survey, 1 muskrat damage site from 2009 was re-evaluated.

Site #	Damage Type in 2009	Damage Type in 2010/ Condition
422	Storm damage (no muskrat visible)	Recovered

Table 2

Conclusions

The 2010 vegetative damage survey yielded a total of 2,260 acres of nutria damage along transect lines. This figure, when extrapolated, demonstrates that 8,475 acres were impacted coast wide at the time of survey. When compared to 2009 (5,422 acres or 20,333 acres extrapolated coast wide), there was approximately a 58% decrease in the number of damaged acres.

Due to the distance between survey lines, all areas impacted by nutria herbivory could not be identified. Additionally, there were survey miles where nutria activity was observed but marsh conditions did not warrant a damage classification. Again, only the most obvious impacted areas were detected so the total impact of nutria was probably underestimated, however the trend in both decreasing damage acreage and increased marsh recovery are significant.

Section 3

CNCP: Summary of Results (2002-2010) and Adaptive Management

Since the beginning of the Coastwide Nutria Control Program, the number of nutria damage sites observed by aerial surveys has continued to decline (Figure 15.)




Figure 15

Three years prior to implementation of CNCP incentive payments.

	Nutria Harvested		Herbivory Damage (acres)
1999-2000	20,110	2000	97,271
2000-2001	29,544	2001	83,021
2001-2002	24,683	2002	79,444

Table 3

Eight years of CNCP incentive payment implementation.

	Nutria Harvested		Herbivory Damage (acres)
2002-2003	308,160	2003	82,080
2003-2004	332,396	2004	63,398
2004-2005	297,535	2005	53,475
2005-2006	168,843	2006	55,755
2006-2007	375,683	2007	34,665
2007-2008	308,212	2008	23,141
2008-2009	334,038	2009	20,333
2009-2010	445,963	2010	8,475

Table 4

Once again proving the Coastwide Nutria Control Program successful, the 2009-2010 season ended with a record harvest as well as fewer impacted acres. To date, nutria harvest in coastal Louisiana has increased to an average of 321,353 animals per year, and the number of damage acres continues to decrease.

As in the past, CNCP applications will be sent to all participants who submitted applications over the last two years. LDWF will also continue the coordination with trappers and fur buyers/dealers to encourage the maximum use of the entire animal, and landowners will be encouraged to trap/hunt the existing damage sites.

Appendix A.
A Comparison of Seasons 1-8
(2002-2010)

PARISH	2002-2003		2003-2004		2004-2005		2005-2006		2006-2007	
	Nutria Harvested	Percentage	Nutria Harvested	Percentage	Nutria Harvested	Percentage	Nutria Harvested	Percentage	Nutria Harvested	Percentage
Ascension	2,710	0.90%	5,474	1.60%	1,858	0.60%	1,678	1.00%	2,226	0.59%
Assumption	3,128	1.00%	814	0.20%	428	0.10%	2,307	1.40%	2,095	0.56%
Calcasieu	143	-	374	0.10%	448	0.20%	58	0.00%	19	0.01%
Cameron	7,851	2.60%	8,701	2.60%	16,617	5.60%	3,744	2.20%	1,725	0.46%
Iberia	1,412	0.50%	1,960	0.60%	3,521	1.20%	3,014	1.80%	18,910	5.03%
Iberville	0	-	1,567	0.50%	5,559	1.90%	2,360	1.40%	9,172	2.44%
Jefferson	20,529	6.70%	24,896	7.50%	11,036	3.70%	2,875	1.70%	10,405	2.77%
Jefferson Davis	121	-	85	-	175	0.10%	110	0.10%	0	0.00%
Lafayette	39	-	25	-	10	0.00%	0	-	0	0.00%
Lafourche	28,852	9.40%	51,736	15.60%	32,411	10.90%	24,668	14.60%	28,038	7.46%
Livingston	2,631	0.90%	357	0.10%	911	0.30%	1,921	1.10%	1,250	0.33%
Orleans	597	0.20%	0	-	538	0.20%	0	-	575	0.15%
Plaquemines	63,208	20.50%	86,720	26.10%	39,043	13.10%	1,816	1.10%	5,815	1.55%
St. Bernard	5,769	1.80%	13,344	4.00%	4,344	1.50%	0	-	291	0.08%
St. Charles	11,169	3.60%	12,672	3.80%	15,867	5.30%	13,807	8.20%	18,690	4.97%
St. James	95	-	487	0.20%	2,841	1.00%	4,912	2.90%	7,111	1.89%
St. John the Baptist	18,450	6.00%	6,137	1.80%	8,404	2.80%	6,384	3.80%	15,786	4.20%
St. Martin	11,425	3.70%	15,039	4.50%	31,656	10.60%	15,903	9.40%	113,629	30.25%
St. Mary	26,004	8.40%	16,277	4.90%	20,940	7.00%	21,023	12.50%	34,693	9.23%
St. Tammany	4,638	1.50%	3,756	1.10%	5,175	1.70%	1,423	0.80%	2,067	0.55%
Tangipahoa	1,245	0.40%	745	0.20%	565	0.20%	826	0.50%	1,843	0.49%
Terrebonne	92,831	30.10%	72,846	21.90%	81,135	27.30%	57,756	34.20%	99,433	26.47%
Vermilion	5,313	1.70%	8,584	2.60%	14,503	4.70%	2,258	1.30%	1,813	0.48%
West Baton Rouge	-	-	-	-	-	-	-	-	97	0.03%
Total	308,160	99.90%	332,596	99.90%	297,535	100.00%	168,843	100.00%	375,683	100.00%

Table 5. Nutria harvested by parish seasons 1-8, Coastwide Nutria Control Program.

PARISH	2007-2008		2008-2009		2009-2010							
	Nutria Harvested	Percentage	Nutria Harvested	Percentage	Nutria Harvested	Percentage						
Ascension	1,957	0.63%	7,029	2.10%	7,049	1.58%						
Assumption	3,863	1.25%	1,093	0.33%	2,930	0.66%						
Calcasieu	19	0.01%	0	0.00%	0	0.00%						
Cameron	649	0.21%	1,245	0.37%	1,177	0.26%						
Iberia	6,119	1.99%	978	0.29%	1,206	0.27%						
Iberville	2,105	0.68%	231	0.07%	6,065	1.36%						
Jefferson	11,299	3.67%	12,515	3.75%	11,506	2.58%						
Jefferson Davis	0	0.00%	0	0.00%	0	0.00%						
Lafayette	0	0.00%	0	0.00%	0	0.00%						
Lafourche	25,473	8.26%	48,252	14.45%	39,564	8.87%						
Livingston	695	0.23%	444	0.13%	2,186	0.49%						
Orleans	1,333	0.43	656	0.20%	1,756	0.39%						
Plaquemines	41,072	13.33%	42,212	12.64%	69,294	15.54%						
St. Bernard	4,150	1.35%	13,965	4.18%	3,543	0.79%						
St. Charles	18,271	5.93%	21,215	6.35%	27,221	6.10%						
St. James	9,604	3.12%	8,990	2.69%	19,226	4.31%						
St. John the Baptist	6,728	2.18%	10,189	3.05%	6,642	1.49%						
St. Martin	54,726	17.76%	44,972	13.46%	63,619	14.27%						
St. Mary	34,210	11.10%	34,811	10.42%	67,631	15.17%						
St. Tammany	4,356	1.41%	5,680	1.70%	8,855	1.99%						
Tangipahoa	2,323	0.75%	4,974	1.49%	267	0.06%						
Terrebonne	78,934	25.61%	74,587	22.33%	106,226	23.82%						
Vermilion	326	0.11%	0	0.00%	0	0.00%						
West Baton Rouge	0	0.00%	0	0.00%	0	0.00%						
Total	308,212	100.00%	334,038	100.00%	445,963	100%						

Table 5 (Continued). Nutria harvested by parish seasons 1-8, Coastwide Nutria Control Program.

PARISH	2002-2003			2003-2004			2004-2005			2005-2006		
	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun
Ascension	0	2,306	404	0	4,093	1,381	100	1,678	80	470	908	300
Assumption	284	2,786	58	47	767	0	188	106	134	1,454	711	143
Calcasieu	0	143	0	0	374	0	213	24	212	57	1	0
Cameron	3,611	4,210	30	4,974	3,639	89	5,779	8,961	1,877	1,362	583	1,799
Iberia	0	1,353	59	636	1,324	0	1,286	1,310	926	1,215	449	1,350
Iberville	0	0	0	717	850	0	4,348	1,211	0	1,156	622	582
Jefferson	5,869	14,094	566	12,991	11,835	70	6,286	4,307	443	2,234	477	164
Jefferson Davis	121	0	0	85	0	0	158	18	0	109	1	0
Lafayette	19	10	10	0	25	0	0	10	0	0	0	0
Lafourche	11,807	16,826	219	28,516	22,780	440	12,221	18,212	1,977	9,113	11,000	4,555
Livingston	0	2,631	0	0	336	21	0	911	0	0	1,921	0
Orleans	287	219	91	0	0	0	538	0	0	0	0	0
Plaquemines	9,899	52,933	376	34,683	51,302	735	18,121	20,642	280	343	843	630
St. Bernard	2,877	2,892	0	5,412	7,783	149	727	3,617	0	0	0	0
St. Charles	2,099	8,706	364	2,801	9,543	329	1,279	13,958	631	1,863	10,915	1,029
St. James	48	47	0	97	350	40	32	2,752	57	278	4,239	395
St. John the Baptist	1,505	11,132	5,813	2,517	2,200	1,420	2,971	4,788	645	2,165	3,488	538
St. Martin	1,497	9,593	335	5,784	8,790	465	10,684	9,703	11,269	4,137	5,355	6,412
St. Mary	11,073	14,849	82	6,616	9,619	42	9,700	10,798	442	9,266	11,202	554
St. Tammany	3,088	1,529	21	2,687	1,069	0	2,692	2,483	0	533	800	90
Tangipahoa	335	894	16	577	169	0	35	530	0	142	638	46
Terrebonne	46,761	45,317	753	44,419	26,335	2,092	31,730	45,893	3,512	28,132	25,577	4,047
Vermilion	2,370	2,729	214	5,119	3,435	30	5,580	7,900	572	1,076	1,182	0
West Baton Rouge	0	0	0	0	0	0	0	0	0	0	0	0
*Total	103,550	195,199	9,411	158,678	166,618	7,303	114,668	159,810	23,057	65,104	80,912	22,634

Table 6. Method of take by parish for seasons 1-8, Coastwide Nutria Control Program

* Totals may not be exact due to reporting of percentages.

PARISH	2006-2007			2007-2008			2008-2009			2009-2010		
	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun	Trap	Rifle	Shot Gun
Ascension	0	2,008	218	0	1,905	52	217	6,751	61	338	6,712	0
Assumption	354	686	1,056	634	2,944	285	85	933	75	546	1,916	469
Calcasieu	19	0	0	19	0	0	0	0	0	0	0	0
Cameron	347	902	477	509	70	70	1,062	128	55	1,177	0	0
Iberia	6,695	4,635	7,580	3,623	1,248	1,247	258	524	196	932	274	0
Iberville	4,907	460	3,860	754	508	843	103	0	128	4,051	1,670	344
Jefferson	4,731	5,568	106	3,901	6,456	943	4,185	8,146	184	3,164	8,202	140
Jefferson Davis	0	0	0	0	0	0	0	0	0	0	0	0
Lafayette	0	0	0	0	0	0	0	0	0	0	0	0
Lafourche	12,279	11,480	4,279	9,702	11,425	4,345	32,373	13,324	2,555	21,796	16,310	1,458
Livingston	0	1,250	0	0	695	0	0	444	0	460	1,726	0
Orleans	575	0	0	1,333	0	0	656	0	0	1,658	71	27
Plaquemines	3,200	2,554	61	30,093	10,609	0	21,394	19,372	1,447	25,379	43,480	436
St. Bernard	146	146	0	4,071	79	370	9,790	4,131	43	3,177	240	126
St. Charles	6,637	9,401	2,652	3,607	13,366	1,298	6,111	14,036	1,068	7,712	18,593	916
St. James	203	6,439	469	425	9,128	51	597	7,862	531	572	17,805	849
St. John the Baptist	4,223	9,215	2,348	2,323	3,834	572	1,490	8,372	327	2,856	3,776	10
St. Martin	39,972	35,737	37,920	27,937	17,123	9,666	21,134	17,512	6,326	43,341	12,952	7,326
St. Mary	12,810	19,997	1,886	10,783	21,304	2,123	13,357	18,480	2,974	13,026	51,170	3,435
St. Tammany	1,452	529	86	1,736	2,216	404	3,377	1,848	456	2,604	4,945	1,307
Tangipahoa	542	1,189	113	563	1,760	0	321	4,530	124	0	267	0
Terrebonne	36,867	51,357	11,209	28,055	45,000	5,879	25,846	46,139	2,602	40,669	62,264	3,292
Vermilion	1,174	494	145	262	65	0	0	0	0	0	0	0
West Baton Rouge	0	97	0	0	0	0	0	0	0	0	0	0
*Total	137,133	164,144	74,465	130,330	149,734	28,148	142,356	172,531	19,151	173,456	252,373	20,134

Table 6 (continued). Method of take by parish for seasons 1-8, Coastwide Nutria Control Program

* Totals may not be exact due to reporting of percentages.

Year	Number of sites surveyed	Number of sites with current damage	Number of site converted to open water	Sites with vegetative recovery
2002	108 ¹	86	8	12
2003	100	81	3	16
2004	93	68	1	24
2005	78	47	2	29
2006	52	31	9	12
2007	34	23	3 (partial sites)	11 ²
2008	23	16	1 (partial site)	6
2009	24	19	1 (partial site)	5 ²
2010	20	11	0	9

Table 7. Status and number of nutria herbivory sites surveyed from 2002 to 2010.

¹ Two sites could not be evaluated due to high water.

² Total includes 1 site with partial recovery.

PARISH	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Terrebonne	41	12,951	34	12,521	27	7,679	18	4,541	14	7,340
Lafourche	8	1,222	7	610	5	381	2	127	0	0
Jefferson	17	3,003	10	1,805	9	1,718	7	1,383	5	874
Plaquemines	10	882	13	2,540	7	2,494	7	1,850	7	1763
St. Charles	6	768	6	1,266	9	2,564	6	4,690	5	3249
Cameron	0	0	0	0	0	0	0	0	1	233
St. Bernard	6	921	5	918	5	1,035	4	882	4	1,004
St. John	0	0	1	20	2	111	2	240	2	241
Iberia	0	0	0	0	0	0	1	158	0	0
St. Tammany	4	752	2	360	0	0	0	0	0	0
Orleans	2	686	2	962	0	0	0	0	0	0
St. Mary	0	0	0	0	0	0	0	0	0	0
Vermilion	0	0	4	886	5	924	2	389	1	76
Jefferson Davis	0	0	0	0	0	0	0	0	1	88
St. John the Baptist	0	0	0	0	0	0	0	0	0	0
Total	94	21,185 ¹	84	21,888 ¹	69	16,906 ¹	49	14,260 ¹	40	14,868 ^{1,2}

Table 8. Number of nutria damaged sites and acres damaged along transects by parish in coastal Louisiana, 2002 - 2009.

¹This figure represents acres damaged along transects only. Actual damage coast wide is approximately 3.75 times larger than the area estimated by this survey.

²This figure includes 2,553 acres of marsh previously impacted by nutria that was likely converted to open water in Plaquemines and St. Bernard Parishes due to tidal scour from Hurricane Katrina.

³These figures include acres from sites that were partially converted to open water.

PARISH	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF			
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES		
Terrebonne	12	5,915	12	3,768	10	3,162	10	2,241		
Lafourche	2	328	2	338	2	207	1	19		
Jefferson	3	177 ³	2	69	1	29	0	0		
Plaquemines	0	0	1	11	1	9	0	0		
St. Charles	4	2,216 ³	5 ³	2,215 ³	4	1,895	0	0		
Cameron	1	167	0	0	1	120	0	0		
St. Bernard	1	225 ³	0	0	0	0	0	0		
St. John	0	0	0	0	0	0	0	0		
Iberia	0	0	0	0	0	0	0	0		
St. Tammany	0	0	0	0	0	0	0	0		
Orleans	0	0	0	0	0	0	0	0		
St. Mary	0	0	0	0	0	0	0	0		
Vermilion	0	0	0	0	0	0	0	0		
Jefferson Davis	1	81	0	0	0	0	0	0		
St. John the Baptist	1	135	1	70	0	0	0	0		
Total	25	9,244 ^{1,3}	23	6,471 ^{1,3}	19	5,422 ¹	11	2,260 ¹		

Table 8 (Continued). Number of nutria damaged sites and acres damaged along transects by parish in coastal Louisiana, 2002 - 2010.

¹This figure represents acres damaged along transects only. Actual damage coast wide is approximately 3.75 times larger than the area estimated by this survey.

²This figure includes 2,553 acres of marsh previously impacted by nutria that was likely converted to open water in Plaquemines and St. Bernard Parishes due to tidal scour from Hurricane Katrina.

³These figures include acres from sites that were partially converted to open water.

MARSH TYPE	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Fresh	41	11,593	36	10,871	37	10,565	26	9,811	23	11,273
Intermediate	39	7,416	31	8,086	25	5,128	19	3,789	16	3,421
Brackish	14	2,176	17	2,931	7	1,213	4	660	1	174
Total	94	21,185	84	21,888	69	16,906	49	14,260	40	14,868

Table 9. Number of nutria damaged sites and acres damaged, by marsh type along transects in coastal Louisiana during 2002 to 2010; number includes sites converted to open water.

¹ Total includes sites that were partially converted to open water.

MARSH TYPE	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF			
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES		
Fresh	21	8,842	21	6,127	17	5,384	11	2,260		
Intermediate	3	298	2	44	2	38	0	0		
Brackish	1	104	0	0	0	0	0	0		
Total	25 ¹	9,244 ¹	23	6,471 ¹	19	5,422	11	2,260		

Table 9 (Continued). Number of nutria damaged sites and acres damaged, by marsh type along transects in coastal Louisiana during 2002 to 2010; number includes sites converted to open water.

¹ Total includes sites that were partially converted to open water.

NUTRIA RELATIVE ABUNDANCE RATING	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
NO NUTRIA SIGN VISIBLE	21	5,990	23	5,972	13	3,569	12	2,992	4	519
NUTRIA SIGN VISIBLE	31	4,379	26	3,562	29	6,040	28	6,748	26	11,223
ABUNDANT FEEDING	17	4,198	19	6,682	19	5,251	4	4,113	1	573
HEAVY FEEDING	17	5,568	14	5,599	7	2,026	1	273	0	0
TOTAL	86	20,135	81	21,815	69	16,886	47	14,126	31	12,315

Table 10. Number of nutria damage sites and acres damaged by revised nutria relative abundance rating in coastal Louisiana during 2002 to 2010; numbers do not include sites converted to open water.

NUTRIA RELATIVE ABUNDANCE RATING	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
NO NUTRIA SIGN VISIBLE	2	73	0	0	0	0	0	0		
NUTRIA SIGN VISIBLE	12	3,402	13	2,234	6	517	0	0		
ABUNDANT FEEDING	5	1,495	8	3,522	8	1,169	7	640		
HEAVY FEEDING	4	3,658	2	415	5	3,736	4	1,620		
TOTAL	23	8,628	23	6,171	19	5,422	11	2,260		

Table 10 (Continued). Number of nutria damage sites and acres damaged by revised nutria relative abundance rating in coastal Louisiana during 2002 to 2010; numbers do not include sites converted to open water.

VEGETATIVE DAMAGE RATING	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
NO VEGETATIVE DAMAGE	1	30	0	0	0	0	0	0	0	0
MINOR VEGETATIVE DAMAGE	28	3,498	26	8,732	35	6,675	34	8,070	21	7,621
MODERATE VEGETATIVE DAMAGE	44	13,156	41	9,221	29	9,536	12	5,905	9	4,581
SEVERE VEGETATIVE DAMAGE	13	3,451	14	3,862	4	675	1	151	1	113
CONVERTED TO OPEN WATER	8	1,050	3	73	1	20	2	134	9	2,553
TOTAL	94	21,185	84	21,888	69	16,906	49	14,260	40	14,868

Table 11. Number of nutria damage sites and number of acres by the vegetative damage rating in coastal Louisiana 2002 to 2010.

¹ Total includes sites that were partially converted to open water.

VEGETATIVE DAMAGE RATING	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
NO VEGETATIVE DAMAGE	0	0	0	0	0	0	0	0		
MINOR VEGETATIVE DAMAGE	17	4,021	17	5,402	15	5,102	11	2,260		
MODERATE VEGETATIVE DAMAGE	6	4,607	5	640	4	320	0	0		
SEVERE VEGETATIVE DAMAGE	0	0	1	129	0	0	0	0		
CONVERTED TO OPEN WATER	3 ¹	616 ¹	1 ¹	300	1 ¹	90	0	0		
TOTAL	26 ¹	9,244 ¹	24 ¹	6,471 ¹	20	5,512	11	2,260		

Table 11 (Continued). Number of nutria damage sites and number of acres by the vegetative damage rating in coastal Louisiana 2002 to 2010.

¹ Total includes sites that were partially converted to open water.

AGE OF DAMAGE AND CONDITON RATING	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Recovered	12	1,119	16	1,674	24	6,049	29	4,169	13 ¹	1,341 ¹
Old Recovering	51	7,694	51	14,382	53	12,338	39	10,878	21	9,429
Old Not Recovering	31	11,449	17	5,375	5	2,898	2	656	4	1,519
Recent Recovering	0	0	0	0	1	35	1	10	0	0
Recent Not Recovering	0	0	0	0	0	0	0	0	1	285
Current Damage	4	992	13	2,058	9	1,615	5	2,582	5	1,082
Total	98	21,254	97	23,489	92	22,935	76	18,295	44 ¹	13,656 ¹

Table 12. Number of nutria damage sites by age of damage and condition rating in coastal Louisiana in 2002 to 2010.

¹ Total includes sites that were partially recovered.

AGE OF DAMAGE AND CONDITON RATING	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Recovered	11 ¹	1,783 ¹	6	736	5 ¹	673 ¹	9	1,914		
Old Recovering	14	5,011	15	3,852	16	5,321	10	2,198		
Old Not Recovering	5	2,874	3	1,914	2	57	0	0		
Recent Recovering	0	0	0	0	0	0	0	0		
Recent Not Recovering	0	0	0	0	0	0	0	0		
Current Damage	4	743	5	405	1	44	1	62		
Total	34 ¹	10,411 ¹	29	6,907	23	6,095	20	4,174		

Table 12 (Continued). Number of nutria damage sites by age of damage and condition rating in coastal Louisiana in 2002 to 2010.

¹Total includes sites that were partially recovered.

PREDICTION OF RECOVERY BY END OF GROWING SEASON	2002		2003		2004		2005		2006	
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Full Recovery	7	919	8	4,238	10	338	6	443	4	828
Partial Recovery	59	13,950	64	14,497	50	13,440	36	10,073	27	11,487
Increased Damage	5	1,086	6	1,646	6	2,811	5	3,610	0	0
No Recovery Predicated	15	4,180	3	1,434	2	297	0	0	0	0
TOTAL	94	21,185	84	21,888	69	16,906	49	14,260	31	12,315

Table 13. Number of nutria damage sites and acres damaged, by prediction of recovery rating in coastal Louisiana in 2002 to 2010.

PREDICTION OF RECOVERY BY END OF GROWING SEASON	2007		2008		2009		2010			
	NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF		NUMBER OF	
	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES	SITES	ACRES
Full Recovery	2	350	1	80	2	1,588	2	84		
Partial Recovery	21	8,278	22	6,091	16	3,543	9	2,176		
Increased Damage	0	0	0	0	1	291	0	0		
No Recovery Predicated	0	0	0	0	0	0	0	0		
TOTAL	23	8,628	23	6,171	19	5,422	11	2,260		


Table 13 (Continued). Number of nutria damage sites and acres damaged, by prediction of recovery rating in coastal Louisiana in 2002 to 2010.

APPENDIX B.
2009 Nutria Vegetative Damage Sites

SITE	MARSH TYPE	LATITUDE	LONGITUDE	DAMAGE TYPE	DAMAGE D ACRES	ACRES TO OPEN WATER	NRAR	VDR	AGE OF DAMAGE	PREDICTION	PARISH
8	F	29.574	-91.17139	Nutria	291	0	3	1	1	2	Terrebonne
9	F	29.5813	-91.12733	Nutria	254	0	1	2	1	2	Terrebonne
17	F	29.5385	-91.04686	Nutria	261	0	2	1	1	2	Terrebonne
60	I	29.7173	-90.04149	Nutria	9	0	1	2	1	2	Jefferson
60B	I	29.716	-90.05147	Nutria	29	0	3	2	2	2	Jefferson
92	F	29.7178	-90.07776	Nutria	36	0	0	0	0	99	Jefferson
94	F	29.8696	-90.2885	Nutria	39	90	0	0	0	99	St. Charles
120	F	29.5907	-91.06539	Nutria	1,457	0	3	1	1	2	Terrebonne
171	F	29.9114	-90.47039	Nutria	1,484	0	3	1	1	3	St. Charles
238	F	29.9272	-90.52978	Hog/Nutria	213	0	2	1	1	2	St. Charles
274	F	29.5649	-91.08909	Nutria	198	0	2	1	1	2	Terrebonne
311	F	29.5514	-90.97915	Nutria	464	0	0	0	0	99	Terrebonne
345	F	29.614	-90.57279	Nutria	80	0	0	0	0	99	Terrebonne
390	F	29.8824	-90.44819	Nutria	104	0	1	1	1	1	St. Charles
400	F	29.5755	-91.11566	Nutria	475	0	3	1	1	2	Terrebonne
418	F	29.5865	-91.01636	Nutria	54	0	0	0	0	99	Terrebonne
419	F	29.6009	-91.01346	Nutria	168	0	2	1	1	2	Terrebonne
420	F	29.6223	-90.64151	Nutria	163	0	2	1	1	1	Lafourche
422	I	29.7318	-92.27	Muskrat/ Storm	152	0	0	0	0	2	Vermillion
423	F	29.5773	-91.19447	Nutria	16	0	1	1	1	2	Terrebonne
424	F	29.485	-91.10953	Nutria	14	0	1	1	1	2	Terrebonne
425	F	29.5588	-91.1008	Nutria	28	0	2	2	2	2	Terrebonne
426	F	29.948	-90.51209	Nutria	94	0	2	1	1	2	St. Charles
427	F	29.9174	-90.62198	Nutria	120	0	1	1	1	2	St. John the Baptist
428	f	29.55822	-90.50727	Nutria	44	0	2	1	5	2	Lafourche

Table 14. 2009 Nutria Vegetative Damage Sites.

**2009-2010
Coastwide Nutria Control Program
Total Tails: 445,963**



APPENDIX C.
Data collected at each damage site during the 2010
vegetative damage survey.

SITE	MARSH TYPE	LATITUDE	LONGITUDE	DAMAGE TYPE	DAMAGE D ACRES	ACRES TO OPEN WATER	NRAR	VDR	AGE OF DAMAGE	PREDICTION	PARISH
8	F	29.56733245	-91.15816697	Nutria	132	0	2	1	1	2	Terrebonne
9	F	29.58074920	-91.11559314	Nutria	65	0	2	1	1	1	Terrebonne
17	F	29.54421508	-91.04159384	Nutria	98	0	2	1	1	2	Terrebonne
60	I	29.7173	-90.04149	Nutria	9	0	0	0	0	99	Terrebonne
60B	I	29.716	-90.05147	Nutria	29	0	0	0	0	99	Terrebonne
120	F	29.59400575	-91.08584744	Nutria	1309	0	3	1	1	2	Terrebonne
171	F	29.9114	-90.47039	Nutria	1484	0	0	0	0	99	Terrebonne
238	F	29.9280	-90.5236	Hog	213	0	2	1	1	2	St. Charles
274	F	29.56536075	-91.08995020	Nutria	206	0	2	1	1	2	Terrebonne
390	F	29.8824	-90.44819	Nutria	104	0	0	0	0	99	Terrebonne
400	F	29.57911833	-91.11267633	Nutria	274	0	3	1	1	2	Terrebonne
418	F	29.58407044	-91.00997214	Nutria	41	0	2	1	1	2	Terrebonne
419	F	29.59461067	-91.01761218	Nutria	36	0	2	1	1	2	Terrebonne
420	F	29.62263639	-90.65223950	Nutria	19	0	3	1	1	1	Lafourche
422	I	29.7318	-92.27	Muskrat/Storm	152	0	0	0	0	99	Terrebonne
423	F	29.5773	-91.19447	Nutria	16	0	0	0	0	99	Terrebonne
424	F	29.485	-91.10953	Nutria	14	0	0	0	0	99	Terrebonne
425	F	29.55648978	-91.10019493	Nutria	18	0	3	1	1	2	Terrebonne
426	F	29.948	-90.51209	Nutria	94	0	0	0	0	99	Terrebonne
427	F	29.9174	-90.62198	Nutria	120	0	0	0	0	99	Terrebonne
428	F	29.55822	-90.50727	Nutria	44	0	0	0	0	99	Terrebonne
429	F	29.54108234	-91.19656897	Nutria	62	0	2	1	5	2	Terrebonne

Table 15. 2010 Nutria Vegetative Damage Sites.

Data Sheet utilized for 2010 nutria herbivory survey.

2010 NUTRIA VEGETATIVE DAMAGE SURVEY

DATE: _____
 TRANSECT#: _____ PHOTOGRAPHY
 MARSH TYPE: _____ FRAME # _____
 LAT: _____ LAT: _____
 LON: _____ LON: _____

LOCATION DESCRIPTION

ON TRANSECT _____
 EAST OF TRANSECT _____
 WEST OF TRANSECT _____ SITE# _____

DAMAGE TYPE

_____ DAMAGE NOT RELATED TO NUTRIA FEEDING
 _____ DAMAGE - STORM RELATED
 _____ DAMAGE - MUSKRAT
 _____ DAMAGE - NUTRIA
 _____ DAMAGE - OTHER _____
 _____ DAMAGED AREA SUBJECT TO TIDAL ACTION: ___ YES ___ NO
 _____ ESTIMATED SIZE OF AREA (ACRES)

NUTRIA RELATIVE ABUNDANCE RATING

VEGETATIVE DAMAGE RATING

_____ NO NUTRIA SIGN VISIBLE (0)	_____ NO VEGETATIVE DAMAGE (0)
_____ NUTRIA SIGN VISIBLE (1)	_____ MINOR VEGETATIVE DAMAGE (1)
_____ ABUNDANT FEEDING (2)	_____ MODERATE VEGETATIVE DAMAGE (2)
_____ HEAVY FEEDING (3)	_____ SEVERE VEGETATIVE DAMAGE (3)
	_____ CONVERTED TO OPEN WATER (4)

NUTRIA VISIBLE IN AREA

_____ WERE NUTRIA SIGHTED: ___ YES ___ NO
 _____ IF YES, HOW MANY? _____

PLANT SPECIES IMPACTED

PLANT SPECIES RECOVERING

PLANT SPECIES ADJACENT

AGE OF DAMAGE AND CONDITION

_____ RECOVERED (0)
 _____ OLD RECOVERING (1)
 _____ OLD NOT RECOVERING (2)
 _____ RECENT RECOVERING (3)
 _____ RECENT NOT RECOVERING (4)
 _____ CURRENT (OCCURRING NOW) (5)

PREDICTION OF RECOVERY BY END OF 2010 GROWING SEASON

_____ NO RECOVERY PREDICTED (0)
 _____ FULL RECOVERY (1)
 _____ PARTIAL RECOVERY (2)
 _____ INCREASED DAMAGE (3) _____ CHECK NEXT YEAR

CODES FOR NUTRIA HERBIVORY SURVEY DATA

¹Marsh Type

Fresh	F
Intermediate	I
Brackish	B

²Nutria Relative Abundance Rating

No Nutria Sign Visible	0
Nutria Sign Visible	1
Abundant Feeding Sign	2
Heavy Feeding	3

³Vegetative Damage Rating

No Vegetative Damage	0
Minor Vegetative Damage	1
Moderate Vegetative Damage	2
Severe Vegetative Damage	3
Converted To Open Water	4

⁴Age of Damage and Condition

Recovered	0
Old Recovering	1
Old Not Recovering	2
Recent Recovering	3
Recent Not Recovering	4
Current (Occurring Now)	5

⁵Prediction of Recovery by End of 2010 Growing Season

No Recovery Predicted	0
Full Recovery	1
Partial Recovery	2
Increased Damage	3

99 – Entry does not apply to this site.